



UNIVERSITI PUTRA MALAYSIA

**DETERMINATION OF THE NUCLEOTIDE SEQUENCE OF
RECOMBINANT PLASMID PBX6 CONTAINING A XYLANASE GENE
FROM *FIBROBACTER SUCCINOGENES* S 85**

YADI SURYADI

FSAS 2001 39

**DETERMINATION OF THE NUCLEOTIDE SEQUENCE OF
RECOMBINANT PLASMID PBX6 CONTAINING A XYLANASE GENE
FROM *FIBROBACTER SUCCINOGENES* S 85**

By

YADI SURYADI

**Thesis Submitted in Fulfilment of the Requirement for the
Degree of Master of Science in the
Faculty of Science and Environmental Studies
Universiti Putra Malaysia**

October 2001



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Science

**DETERMINATION OF THE NUCLEOTIDE SEQUENCE OF
RECOMBINANT PLASMID PBX6 CONTAINING A XYLANASE GENE
FROM *FIBROBACTER SUCCINOGENES* S 85**

By

YADI SURYADI

October 2001

Chairman: Associate Professor Abdullah Sipat, Ph.D.

Faculty: Science and Environmental Studies

pBX6 is a recombinant plasmid containing a 3 kb fragment of insert DNA from *Fibrobacter succinogenes* S 85 genomic DNA which encodes xylanase activity. The recombinant plasmid pBX6 has shown xylanase activity when grown on RBB-xylan plates supplemented with ampicillin. Restriction mapping of the plasmid established the presence of single sites for *EcoRI*, *KpnI*, *Apal*, *SacI*, *EcoRV* and *PstI*. This work was focussed on nucleotide sequence determination of recombinant plasmid pBX6 containing a xylanase gene from *F. succinogenes* S 85. The primer walking strategy was performed to complete manual sequencing, following the protocol cycle sequencing of Sanger dideoxy chain termination. Custom synthesis primer was designed and constructed from 3' end of the preceding sequence on both strands.

The complete nucleotide sequence of plasmid pBX6 has been obtained which overlapped in complementary region. A total of nucleotide sequence of *Pst*I-*Eco*RI DNA fragment has been determined containing bases 3227. Analysis of the whole nucleotide sequence revealed two putative Open Reading Frames (ORFs) i.e; ORF 1 and ORF 2. These ORFs were correspond with other microbial sequences encoding xylanase gene. The ORF 1 starts from bases 129 and ended at 1364, while ORF2 stretches from 1776 to 3227 bases of the insert DNA. Each of the ORF was preceded by putative *E. coli* -10 and -35 promoters. The ORF 1 was translated into a peptide of 411 amino acid residues, whereas ORF2 translated into 483 amino acid residues, respectively. It was suggested that ORF 2 containing xylanase gene was slightly similar with *xynC* gene of *F. succinogenes* S 85.

On the basis of amino acid sequence similarity on both ORF 1 and ORF 2, it was shown that the gene involved in the xylan degradation is closely related to the component family 10 catalytic domains of glycosyl hydrolase. The translated product of amino acid residues was expected to have molecular mass about 45.6 kDa for ORF1 and 52.5 kDa for ORF2, with theoretical pI 4.73 and 5.26, respectively. The highest amino acid composition was glycine, whilst the lowest was cystine. Total number of amino acids which showed negatively charged residues (aspartate-glutamate) was 117 residues, whereas total number of positively charged residues (arginine-lysine) was 86 amino acids. The strongest hydrophobic

region was observed at residues 255-256 amino acid of ORF1, and at 17-18 amino acid residues of ORF2. There were found 7 cysteine residues and containing almost an equal amount of α - β helical structure in the sequence, that increase stability of the protein product. The translated product also showed similarity with other prokaryotic xylanases, which have shown putative protein product ranging from 42% to 92%. The ORF 1 residue of xylanase from *F. succinogenes* S 85 has showed homology with other bacterial xylanases such as *Pseudomonas fluorescens* and *Butyrivibrio fibrisolvens*, ranging from 57% to 94% while ORF 2 cluster together and showed homology ranging from 88% to 92% with xylanase from *F. succinogenes* S 85 (*xynC*, *cmc-xyl*).

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENENTUAN JUJUKAN NUKLEOTIDA PLASMID REKOMBINAN PBX6
YANG MENGANDUNGI GEN XILANASE DARI
FIBROBACTER SUCCINOGENES S 85**

Oleh

YADI SURYADI

Oktober 2001

Pengerusi: Profesor Madya Abdullah Sipat, Ph.D.

Fakulti: Sains dan Pengajian Alam Sekitar

pBX6 ialah plasmid rekombinan yang mengandungi satu DNA selitan dari DNA genomik *F. succinogenes* S 85 yang mengkodkan aktiviti xilanase. Plasmid rekombinan pBX6 telah menunjukkan aktiviti xilanase apabila ditumbuhkan pada petri RBB-xilan yang ditambahkan dengan ampisilin. Pemetaan pembatas bagi plasmid telah mengasaskan kehadiran tapak tunggal untuk *EcoRI*, *KpnI*, *Apal*, *SacI*, *EcoRV* dan *PstI*. Kajian ini menumpukan perhatian pada penentuan jujukan nukleotida plasmid rekombinan pBX6 yang mengandungi gen xilanase dari *F. succinogenes* S 85. Strategi 'primer walking' telah dijalankan untuk melengkapkan jujukan pBX6 secara manual dengan mengikuti kaedah jujukan rangkaian bagi 'pengakhiran rantai dideosi Sanger'. Primer sintesis telah dirangka dan dibina daripada 3' hujung dari jujukan pendahuluan pada kedua-dua rantai.

Jujukan nukleotida penuh bagi potongan DNA *PstI-EcoRI* telah ditentukan mempunyai 3227 bes. Penganalisaan pada seluruh jujukan nukleotida

memberikan 2 anggapan rangka bacaan terbuka (RBT) iaitu RBT 1 dan RBT 2. RBT ini berkenaan rapat dengan jujukan pelbagai mikroorganisma yang mengkodkan gen xilanase. RBT 1 bermula dari bes 129 dan berakhir pada 1364, manakala RBT 2 bermula dari 1776 dan berakhir pada 3227 bes daripada DNA selitan. Setiap RBT didahului jangkakan promoter *E. coli* -10 dan -35. Masing-masing RBT 1 telah ditranslasi kepada satu peptid yang mengandungi 411 asid amino, manakala RBT 2 telah ditranslasi kepada 483 asid amino. Dicaadangkan bahawa ORF 2 mengandungi gen xilanase hampir sama dengan gen *xynC* dari *F. succinogenes* S 85

Berdasarkan kehampirsamaan jujukan asid amino gen xilanase dari *F. succinogenes* S 85 pada kedua RBT 1 dan RBT 2, ditunjukkan bahawa gen yang terlibat pada penguraian xilan berkenaan rapat dengan kumpulan domain katalitik keluarga 10 glikosil hidrolase. Secara teoritikal, produk translasi dari baki asid amino dijangka mempunyai jisim molekul sebanyak 45.6 kDa untuk RBT 1 dan 52.5 kDa untuk RBT 2, dengan masing-masing mempunyai pI 4.73 dan 5.26. Kandungan asid amino yang paling banyak ialah glisina, manakala yang paling sedikit ialah sistina. Jumlah baki asid amino yang bercas negatif (asid glutamat-asid aspartat) ialah 117, manakala jumlah baki yang bercas positif ialah 86 asid amino. Kawasan hidrofobik yang paling kuat telah dijumpai pada asid amino 255-256 bagi RBT 1, dan pada asid amino 17-18 bagi RBT 2. Terdapat 7 baki sistina dan mengandungi hampir sama jumlah struktur helikal α - β dalam jujukannya, ini meninggikan kestabilan produk protein. Produk protein juga menunjukkan

homologi dengan xilanase dari prokariot lainnya dan dijangka mempunyai kesamaan 42% sampai 92%. Produk RBT 1 menunjukkan homologi dengan xilanase bakteri yang lain seperti *Pseudomonas fluorescense* dan *Butyrivibrio fibrisolvens* (57% sampai 94%), manakala RBT 2 menunjukan kesamaaan dengan xilanase dari *F. succinogenes* S 85 (*xynC*, *cmc- xyl*) (88% sampai 92%).

ACKNOWLEDGEMENTS

Alhamdulillah, praise be to ALLAH, the most merciful and the most compassionate for giving me strength and comfort upon completion this research project and thesis.

I would like to express my deepest sincere thank to my chairperson Associate Professor Dr. Abdullah Sipat, Department of Biochemistry and Microbiology, Faculty of Science and Environmental Studies University Putra Malaysia for his excellent guidance, encouragement and constructive suggestions throughout undertaking of my study.

I am grateful to my supervisory committee members, Associate Professor Dr. Khatijah Mohd. Yusoff and Dr. Hirzun Mohd. Yusoff for their support and discussions.

I am indebted to the government of Indonesia, especially to Agriculture Research Management (ARM-II) Project and Director of Research Institute for Food Crops Biotechnology (RIFCB) for their financial support and allowing me to do a postgraduate study in Malaysia. I also wish to thank the Government of Malaysia (Ministry of Science, Technology and Environment) for funding IRPA project of thermostable xylanase gene.

Thanks are due to my lab mates and colleagues in lab. 202 and 143 especially Ainu, Ong, Filza, Yusuf, Aneela, Gayah, Sieh Hie, Wong, Lau, Pria, Becky, Wawa, Dela, and Aini for their friendship and help which make my stay in Malaysia pleasant and had a wonderful time.

Many thanks are devoted to my supportive parents, parents-in law and all family members for all support and guidance. I dedicate this thesis to my beloved wife Lilis Sofiarsih, and daughters Nandina and Rininta, thank you all for your understanding, support and patience throughout my stay abroad.

I certify that an Examination Committee met on 24th October to conduct the final examination of Yadi Suryadi, on his Master of Science thesis entitled "Determination of the Nucleotide Sequence of Recombinant Plasmid pBX6 Containing a Xylanase Gene from *Fibrobacter succinogenes* S85" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The committee recommends that the candidate be awarded the relevant degree. Members of the examination are as follows:

RAJA NOOR ZALIHA RAJA A. RAHMAN, Ph.D,
Associate Professor,
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Chairman)

ABDULLAH SIPAT, Ph.D,
Associate Professor,
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

KHATIJAH MOHD. YUSOFF, Ph.D,
Associate Professor,
Faculty of Science and Environmental Studies
Universiti Putra Malaysia
(Member)

HIRZUN MOHD. YUSOFF, Ph.D,
Fermentation Technology Centre
Institute of Bioscience
Universiti Putra Malaysia
(Member)



MOHD. GHAZALI MOHAYIDIN, Ph.D.
Professor/Deputy Dean of Graduate
School
Universiti Putra Malaysia

Date: 23 NOV 2001

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science.



AINI IDERIS, Ph.D.
Professor/Dean of Graduate
School
Universiti Putra Malaysia

Date:

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



YADI SURYADI

Date: 24th October 2001

TABLE OF CONTENTS

	Page
ABSTRACT.....	ii
ABSTRAK.....	v
ACKNOWLEDGEMENTS.....	viii
APPROVAL	x
DECLARATION.....	xii
LIST OF TABLES.....	xv
LIST OF FIGURES.....	xvi
LIST OF PLATES	xviii
LIST OF ABBREVIATIONS	xix
CHAPTER	
I INTRODUCTION.....	1
II LITERATURE REVIEW	4
Xylan and its Chemical Structure	4
Xylanase	5
Microorganisms Producing Xylanase.....	7
Molecular Studies on Xylanase Gene.....	9
Cloning of Xylanase Gene.....	9
Sequencing of Xylanase Gene.....	13
Strategy for Nucleotide Sequence Determination.....	17
Primer Design and Construction.....	21
Bioinformatics and Sequence Analysis Tools.....	22
Search for Gene	25
Nucleotide Sequence of Xylanase Gen.....	27
Objectives.....	29
III MATERIALS AND METHODS	30
Xylanolytic Plasmid pBX6 Properties	30
General Procedure, Source of Plasmid and Culture Condition	30
Crude Xylanolytic Activity, SDS-PAGE and Dot Blot ELISA..	30
Nucleotide Sequencing	35
Plasmid Extraction, Purification and Restriction Endonuclease	
Analysis.....	35
Primer Walking Strategy, Design and Construction.....	37
Template Preparation and Cycle Sequencing.....	37
Preparation of Gel Casting, Gel loading and Electrophoresis	
Condition.....	41
Autoradiography.....	42

Deletion Mutation of Xylanolytic pBX6 Using PCR.....	42
PCR Condition.....	42
Subcloning of PCR Product.....	43
Data Analysis.....	47
IV RESULTS AND DISCUSSION.....	48
Xylanolytic Plasmid PBX6 and Its Properties.....	48
Nucleotide Sequencing.....	57
Restriction Enzyme Analysis of pBX6.....	57
Primer Walking to Sequence <i>Pst</i> I- <i>Eco</i> RI DNA Fragment.....	60
Restriction Enzyme Analysis.....	64
Prediction of the Putative ORF 1 and ORF 2 for the Xylanase Gene	64
Comparison of ORF 1 and ORF 2 Deduced Sequence to Other Xylanase Genes.....	71
Homology Analysis	71
Comparison between ORF 1 and ORF 2 Sequence.....	73
Phylogenetic tree for the Putative ORF 1 and ORF 2.....	77
Analysis of Characteristic of the Putative ORF 1 and ORF2	83
Amino Acid Composition.....	83
Hydrophobicity and Stability of the Enzyme.....	88
Secondary Structure.....	94
Deletion Mutation of pBX6 Using PCR.....	97
V CONCLUSION.....	104
REFERENCES	107
APPENDICES.....	115
VITAE.....	125

LIST OF TABLES

Table	Page
1 Various recombinant plasmid from <i>F. succinogenes</i> S 85 that express xylanase activity on <i>E. coli</i> HB 101	10
2 Oligonucleotide primers used in sequencing pBX6 based upon primer walking strategy	39
3 Amino acid composition of ORF 1 and ORF 2 from <i>F. succinogenes</i> S 85 encoding xylanase gene.....	86
4 Amino acid composition of xylanase from different prokaryotes.....	87

LIST OF FIGURES

Figure		Page
1	General Structure of Xylan	6
2	Restriction Map of Recombinant Plasmid pBX6 in pUC19 Vector Containing DNA Fragment Insert of 3 kb <i>Pst</i> I- <i>Eco</i> RI.....	12
3	Cycle Sequencing by Chain Dideoxy Termination Method ...	15
4	Autoradiograph of DNA Sequencing Gel.....	16
5	Scheme of Primer Walking Strategy	19
6	Bioinformatics Flow Chart in Nucleotide Sequence	24
7	Flow Chart for the Preparation of Crude Xylanase Enzyme Obtained from pBX6 by Centrifugation	33
8	Schematic Representation of the Primer Walking Strategy for DNA Sequencing	38
9	Deletion Mutation of pBX6 by PCR	46
10	Temperature- Activity Profile of Crude Xylanase	51
11	pH- Activity Profile of Crude Xylanase	52
12	Nucleotide Sequence of pBX6 from <i>F. succinogenes</i> S 85 Coding for Xylanase Gene and its Flanking Regions.....	63
13	Restriction Enzyme Map of the Total of Nucleotide Sequence of the Insert DNA	65
14	Scheme Representing the Putative Xylanase Gene on pBX6 Sequence	68
15	Similarity between Xylanase Genes in ORF 1 and ORF 2 of pBX6 Sequence.....	74
16	Domain Structure of Family Glycosyl Hydrolase F 10 in Amino Acid Sequence of pBX6 Encoding Xylanase from <i>F. succinogenes</i> S 85 (ORF 1 and ORF 2) based upon Pfam Analysis.....	75

17a	Homology of <i>F. succinogenes</i> S 85 (ORF1) Against Several Xylanase Family F-10 Glycosyl Hydrolase.....	78
17b	Homology of <i>F. succinogenes</i> S 85 (ORF2) Against Several Xylanase Family F-10 Glycosyl Hydrolase.....	79
18a	Dendrogram Similarity between Xylanase Gene from <i>F. succinogenes</i> S 85 (ORF 1) with Different Prokaryotes Using Phylogenetic Distance Analysis.....	81
18b	Dendrogram Similarity between Xylanase Gene from <i>F. succinogenes</i> S 85 (ORF 2) with Different Prokaryotes Using Phylogenetic Distance Analysis.....	82
19a	Transmembrane Analysis of ORF 1 Regions.....	90
19b	Transmembrane Analysis of ORF 2 Regions.....	91
20	Hydrophobicity Profile of ORF 1 and ORF 2 Encoding Xylanase Gene from <i>F. succinogenes</i> S 85	92
21	Secondary Structure Prediction of ORF 1 and ORF 2 Amino Acid Residues.....	95
22	PCR Amplification Product Using Primer Combination Fa37/F2Y for Deletion Analysis.....	99

LIST OF PLATES

Plate		Page
1	Screening of Xylanase Activity of Recombinant <i>E. coli</i> HB 101 (pBX6) on 2 YT + RBB-Xylan (0.2%) + Ampicillin.....	50
2	SDS-PAGE of Polyclonal Antibody Derived from Crude pBX6	54
3	Dot blot ELISA Using pAb to Detect Crude Filtrate of Xylanase	56
4	Agarose Electrophoresis of Plasmid pBX6.....	58
5	Restriction Digest of pBX6 Using Different Restriction Enzymes	59
6	A portion of Sequencing Gel (at 660 Position in the Gene) Detected by Autoradiography.....	70
7	PCR Product of pBX 6 Using Primer Pair Combination	100
8	Screening of Recombinant Colony after TOPO TA Cloning to PCR Product (Fa37/F2Y).....	102
9	Clearing Zone of Recombinant Colony from PCR Product (Fa37/F2Y).....	102
10	PCR Confirmation Test for Fragment PCR Product (Fa37/F2Y).....	103

LIST OF ABBREVIATIONS

A	adenine
bp	base pair
C	cytosine
°C	degrees centigrade
DNA	deoxyribonucleic acid
EDTA	ethylene diamine tetra acetic acid
ELISA	enzyme linked immunosorbent assay
G	guanine
<i>g</i>	gravity force
g	gram
h	hour
kb	kilo base pair
kDa	kilo Dalton
M	molar
MCS	multi cloning sites
mA	milliampere
mg	milligram
min	minute
ml	milliliter
mM	millimolar
MW	molecular weight
ng	nanogram

ORF	open reading frame
PCR	polymerase chain reaction
pg	pico gram
pI	isoelectric point
RNA	ribonucleic acid
RBB	remazol brilliant blue
RBS	ribosome binding site
rpm	revolutions per minute
SD	Shine-Dalgarno
SDS	sodium dodecyl sulphate
s	second
TAE	tris acetate- EDTA
TEMED N,N,N,N'	tetramethyl-ethylene diamine
T	thymine
T _m	temperature of melting point
U	unit of xylanase activity
μg	micro gram
μl	micro liter
μCi	micro Curie
YT	yeast tryptone

CHAPTER I

INTRODUCTION

Xylans are heterogeneous polysaccharides found in the cell walls of many plants species (Wong *et al.*, 1986). They are important in their structural roles, being applied as polymers in biomass conversion and other uses (Fry, 1986; Mishra *et al.*, 1989; Wong and Saddler, 1992). Complete breakdown of a branched xylan requires the action of endo β -1,4-xylanase that attacks the polysaccharide backbone and exo β -1,4-xylosidase that hydrolyses xylo-oligosaccharides to xylose (Gilkes *et al.*, 1991). They play an important role in physiological plant tissue and plant defense mechanisms (Esteban *et al.*, 1982). In addition, xylanases are also used in the food and feed industry (Wong *et al.*, 1988).

Enzyme hydrolysis using microbial xylanase is most efficient. Several enzymes, which are capable of degrading xylans, have been isolated from bacteria, yeast and fungi (Gilkes *et al.*, 1991). Although extensive work has been carried out on the purification and characterization of xylanases, more studies at molecular level using recombinant DNA techniques have only been carried out recently (Coughland and Hazlewood, 1993). The rumen bacterium, *Fibrobacter succinogenes* subsp. *succinogenes* S 85 (Montgomery *et al.*, 1988) (previous name: *Bacteroides succinogenes*), is one of the best xylanase producers. Its xylanolytic complex has been investigated (Sipat *et al.*, 1987; Malburg *et al.*, 1993; Ozcan *et al.*, 1996).

Sipat *et al.*, (1987) cloned a 9.4 kb gene fragment into pBR322 from a genomic library of *F. succinogenes* S 85. This recombinant plasmid pBX1 expressed xylanase activity in *Escherichia coli* HB 101. Matte and Forsberg (1992) purified two endoxylanases from *F. succinogenes*, which differed in substrate specificity from the enzyme coded for xylanase gene cloned by Sipat *et al.*, (1987). Malburg *et al.*, (1995) has demonstrated that multiple xylanase genes were possessed by *F. succinogenes* S 85. Coughland and Hazlewood (1993) have also shown that multiple xylanase genes are produced by a large number of hemicellulolytic fungi.

Studies on xylanase function and structure may be useful in determining the efficiency of microbial xylanase. It comprises various combinations of functional elements; such as catalytic domains, cellulose binding domains, linker regions and repeated sequences of amino acids. It is possible to group xylanases into families according to conserved amino acids sequences in the catalytic domains (Henrissat and Bairoch, 1993). All of the xylanases so far, have been characterized which fall into two classes; xylanases, which have high relative molecular mass-low pI class, and xylanases, which have low molecular mass-high pI class (Wong *et al.*, 1988; Henrissat and Bairoch, 1993; Iyo and Forsberg, 1996).

More detailed study of recombinant plasmid derived from *F. succinogenes* S85 carrying xylanase gene is still necessary to increase its efficiency in the industrial application. Molecular aspects such as cloning and sequencing

are useful tools to obtain the nucleotide sequence involved in xylanase gene product, hence allowing manipulation of the enzyme for various purposes (Malburg *et al.*, 1993).

Recombinant plasmid pBX6 consists of the vector pUC19 with a 3 kb DNA insert from *F. succinogenes* S 85 that was previously cloned by Sipat *et al.*, (1987). This thesis reports the study and characterization of plasmid pBX6 encoding xylanase activity from *F. succinogenes* S 85 by determining the nucleotide sequence and its deduced amino acid sequence in order to understand structure and function of xylanase gene.

CHAPTER II

LITERATURE REVIEW

Xylan and its Chemical Structure

Hemicellulose polysaccharides are the major structure component in the plant cell walls which represent an important energy source for the host ruminant (Dekker and Richards, 1976; Williams and Withers, 1992). Hemicelluloses are generally a mixture of xylans with small amount of various mannans, which can be obtained by alkaline extraction of forages and woody biomass (Hespell *et al.*, 1987).

Xylans are generally represented as complex heteropolysaccharides whose structure varies in nature and degree of branching β -1,4-linked xylopyranosyl main chain. They constitute at least 30% of dry weight in plant cell walls. The general features of the xylans are given in Figure 1. The typical of xylans may derived from various structural type; linear homoxylan (no side chain at all), arabinoxylan (has only one side chain), glucuronoxylan (has α D glucuronic acid or 4-O-methyl ether derivative) and glucuronoarabinoxylan (has α L arabinose, α D glucuronic acid and 4-O-methyl α D glucuronic) (Josselau *et al.*, 1992). Xylans may interact with cellulose, other polysaccharides, and lignin hydroxycinnamic acid through various covalent linkages and non-covalent secondary forces (Ethier *et al.*, 1994). The end product of xylan hydrolysis using alkaline or enzyme, can